

**Exam 2020/2021**

October 28, 2020, from 13:45 to 16:45  
 Documents allowed, Internet not allowed  
 Do what you can, and do not worry

$(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$  is a filtered probability space, with complete and right continuous filtration.  
 $B = (B_t)_{t \geq 0}$  is a  $d$ -dimensional Brownian motion issued from the origin,  $d \geq 1$ .

**Exercise 1** (Representation of a process). Take  $d = 1$  and  $x \in \mathbb{R}$ .

- Recall the computations and reasoning showing that the process  $(Z_t)_{t \geq 0}$  defined by

$$Z_t = xe^{-t} + e^{-t}M_t \quad \text{where} \quad M_t = \sqrt{2} \int_0^t e^s dB_s$$

is the unique solution of the stochastic differential equation  $Z_0 = x, dZ_t = \sqrt{2}dB_t - Z_t dt$ .

- Show that for all  $t \geq 0, Z_t \stackrel{\text{law}}{=} xe^{-t} + e^{-t}B_{e^{2t}-1}$ .
- Can we have, for all  $t \geq 0, Z_t = xe^{-t} + e^{-t}B_{e^{2t}-1}$ ?
- Show that the process  $(M_t)_{t \geq 0}$  is a continuous local martingale with, for all  $t \geq 0, \langle M \rangle_t = e^{2t} - 1$ .
- Deduce that there exists a Brownian motion  $(W_t)_{t \geq 0}$  such that for all  $t \geq 0, Z_t = xe^{-t} + e^{-t}W_{e^{2t}-1}$ .

**Exercise 2** (Study of a special process). Let  $d = 1, \alpha \geq 0, x \geq 0$ . Let  $X$  be a continuous semi-martingale taking values in  $\mathbb{R}_+$  and solving the stochastic differential equation:

$$X_t = x + 2 \int_0^t \sqrt{X_s} dB_s + \alpha t, \quad t \geq 0.$$

Let  $f : [0, +\infty) \rightarrow [0, +\infty)$  be continuous and  $\varphi : [0, +\infty) \rightarrow (0, +\infty)$  be positive and  $\mathcal{C}^2$ , solving the ordinary differential equation  $\varphi'' = 2f\varphi$  with boundary conditions  $\varphi(0) = 1$  and  $\varphi'(1) = 0$ . Note that  $\varphi > 0$ .

- Could you give an explicit example of process  $X$  for special values of  $\alpha$ ?
- Show that  $\varphi$  decreases on the interval  $[0, 1]$
- Show that  $u = \varphi'/(2\varphi)$  solves the differential equation  $u' + 2u^2 = f$
- Show that for all  $t \geq 0$ ,

$$u(t)X_t - \int_0^t f(s)X_s ds = u(0)x + \int_0^t u(s)dX_s - 2 \int_0^t u(s)^2 X_s ds.$$

- For all  $t \geq 0$ , let us define  $Y_t = u(t)X_t - \int_0^t f(s)X_s ds$ . Show that

$$\varphi(t)^{-\frac{\alpha}{2}} e^{Y_t} = e^{N_t - \frac{1}{2}\langle N \rangle_t} \quad \text{where} \quad N_t = u(0)x + 2 \int_0^t u(s)\sqrt{X_s} dB_s$$

- Show that

$$\mathbb{E} \exp\left(-\int_0^1 f(s)X_s ds\right) = \varphi(1)^{\frac{\alpha}{2}} e^{\frac{\alpha}{2}\varphi'(0)}$$

- From now on, let  $\lambda > 0$ . Prove that

$$\mathbb{E} \exp\left(-\lambda \int_0^1 X_s ds\right) = (\cosh(\sqrt{2\lambda}))^{-\frac{\alpha}{2}} e^{-\frac{\alpha}{2}\sqrt{2\lambda} \tanh \sqrt{2\lambda}}$$

- Prove that for all  $\lambda > 0$  and  $y \in \mathbb{R}$ ,

$$\mathbb{E} \exp\left(-\lambda \int_0^1 (y + B_s)^2 ds\right) = (\cosh(\sqrt{2\lambda}))^{-\frac{1}{2}} e^{-\frac{y^2}{2}\sqrt{2\lambda} \tanh \sqrt{2\lambda}}$$

**Exercise 3** (Strict local martingales). We take  $d = 3$ ,  $X = x + B$ ,  $0 < r < |x| < R < \infty$ , and, for all  $a \geq 0$ ,

$$T_a = \inf\{t \geq 0 : |X_t| = a\}.$$

1. Show that if  $M = (M_t)_{t \geq 0}$  is a continuous local martingale with for all  $t \geq 0$ ,  $|M_t| \leq U$  where  $U \in L^1$ , then  $M$  is a martingale. Does it remain true if the domination condition is replaced by “ $M$  is u.i.”?
2. Show that if  $Z = (Z_t)_{t \geq 0}$  is  $d$ -dimensional, adapted, taking values in an open set  $D \subset \mathbb{R}^d$ , such that its components are continuous local martingales, and for all  $1 \leq j, k \leq d$ ,  $\langle Z^j, Z^k \rangle = V \mathbf{1}_{j=k}$  for a finite variation process  $V$ , then, for all harmonic  $u : D \rightarrow \mathbb{R}$ , the process  $u(Z)$  is a local martingale.
3. Show that  $|\bullet|^{-1}$  is harmonic on  $\mathbb{R}^3 \setminus \{0\}$ .
4. Show that  $T_R < \infty$  almost surely and

$$\mathbb{P}(T_r < T_R) = \frac{R^{-1} - |x|^{-1}}{R^{-1} - r^{-1}}.$$

5. Deduce from the previous formula that a.s. for all  $t \geq 0$ ,  $X_t \neq 0$ .
6. Show that a.s.  $\lim_{t \rightarrow \infty} |B_t| = +\infty$ . Hint: show that  $|X|^{-1}$  is a non-negative super-martingale.
7. Show that  $|X|^{-1}$  is bounded in  $L^2$ . Hint: density of  $B_t$  in spherical coordinates.
8. Show that  $|X|^{-1}$  is a continuous local martingale, but is not a martingale.

**Exercise 4** (Strict local martingales and stochastic integrals).

1. Give an example of an Itô stochastic integral which is a local martingale but not a martingale, without using the previous exercise.

– oOo –